

ASSIGNMENT 2

POINTS: 50

DATE GIVEN: 01-SEP-2023

DUE: 18-SEP-2023

Rules:

- You are strongly encouraged to work *independently*. That is the best way to understand & master the subject.
- Write the solutions on your own and honorably *acknowledge* the sources if any. <http://cse.iitk.ac.in/pages/AntiCheatingPolicy.html>
- Submit your solutions, before time, to your TA. Please give your LaTeXed or Word processed solution-sheet in PDF. This will be graded, and commented, in-place.
- Clearly express the fundamental *idea* of your proof/ algorithm before going into the other proof details. The distribution of partial marks is according to the proof steps.
- There will be a penalty if you write unnecessary or unrelated details in your solution. Also, do not repeat the proofs done in the class.
- Problems marked '0 points' are for practice.
- Acknowledgements: Several problems are from *Arora & Barak, Computational Complexity: A Modern Approach*, and other lecture notes.

Question 1: [6+3 points] In the class we proved the Razborov-Smolensky result: $\text{mod}_p \notin \text{ACC}^0[q]$, for $p = 2 = q - 1$. Give the general proof for primes $p \neq q$.

What happens to the proof when one of them is a composite?

Question 2: [7 points] Show that the OR of n variables cannot be expressed as a polynomial over \mathbb{F}_p of degree less than n .

Question 3: [13 points] Give an example of a set-family \mathcal{Z} with $(p-1)^\ell$ subsets (each size $\leq \ell$), such that there is no sunflower with p -petals.

Question 4: [6+9 points] Let A be a real symmetric stochastic matrix. Show that

- (1) its eigenvalues are real with absolute value at most 1.
- (2) its eigenvectors, for different eigenvalues, are linearly independent.

Question 5: [6 points] Let A, B be real symmetric stochastic matrices. Show that the *second-largest eigenvalue* function $\lambda(\cdot)$ is *subadditive* (i.e. $\lambda(A + B) \leq \lambda(A) + \lambda(B)$).

Question 6: [0 points] Let G be a graph with only *two* distinct eigenvalues. Prove that G is a *complete graph*.

Question 7: [0 points] Show that for an $n \times n$ matrix $X = ((x_{i,j}))$, the polynomial $\text{per}(X)$ has a formula of size $2^{O(n)}$.

Question 8: [0 points] Show that $\text{AM}[k] = \text{AM}[2]$ for all constant $k \geq 2$.

Question 9: [0 points] Show that the graph isomorphism problem is in $\text{NP} \cap \text{coAM}$. (So, “almost” in $\text{NP} \cap \text{coNP}$.)

Question 10: [0 points] Fix parameters n, d . Show that a random n -vertex d -regular graph is a $(n, 2d, 1/3)$ -combinatorial expander.

Question 11: [0 points] Consider a subset A of the n -tuples \mathbb{F}_3^n . We are interested in the largest set A that is *free of arithmetic progressions of size 3* (i.e. there is no $a, b, c \in A$ satisfying $a + b + c \equiv \mathbf{0} \pmod{3}$).

Could you relate this question to the *sunflower lemma*?

Question 12: [0 points] Recall our proof of the $n^{\Omega(\sqrt{k})}$ monotone circuit lower bound for the k -clique problem. Is this bound tight?

Question 13: [0 points] Think of another “hard” problem that is monotone.

□□□