

CS 771A: Intro to Machine Learning, IIT Kanpur			Quiz I (1 Feb 2022)
Name			20 marks
Roll No		Dept.	Page 1 of 2



**Instructions:**

1. This question paper contains 1 page (2 sides of paper). Please verify.
2. Write your name, roll number, department above in **block letters neatly with ink**.
3. Write your final answers neatly **with a blue/black pen**. Pencil marks may get smudged.
4. Don't overwrite/scratch answers especially in MCQ – such cases will get straight 0 marks.
5. Do not rush to fill in answers. You have enough time to solve this quiz.

**Q1. Write T or F for True/False (write only in the box on the right-hand side) (6 x 1 = 6 marks)**

1	If $\mathcal{P}, \mathcal{Q} \subset \mathbb{R}^2$ are two non-empty convex sets, then their union i.e., $\mathcal{P} \cup \mathcal{Q}$ can never be convex.	
2	$f: \mathbb{R} \rightarrow \mathbb{R}$ is convex, non-differentiable and $-1 \in \partial f(x_0)$ (i.e., -1 is a subgradient), then $f$ must be decreasing around $x_0$ i.e., for some $\epsilon > 0$ , $f(x_0 - \epsilon) > f(x_0 + \epsilon)$ .	
3	If $\mathcal{C} \stackrel{\text{def}}{=} \{(x, y) \in \mathbb{R}^2: 2 \leq x + y \leq 3\}$ is the set of all 2D vectors whose sum of coordinates is between 2 and 3, then $\mathcal{C}$ is a convex set.	
4	If $\mathbf{u}, \mathbf{v} \in \mathbb{R}^3$ are two vectors such that $\ \mathbf{u}\ _2 \leq 1$ and $\ \mathbf{v}\ _2 < 2$ then it is possible that $\mathbf{u}^T \mathbf{v} > 2.5$ for certain values of $\mathbf{u}, \mathbf{v}$ .	
5	If $\mathbf{u}, \mathbf{v} \in \mathbb{R}^3$ are two vectors such that $\ \mathbf{u}\ _2 \leq 1$ and $\ \mathbf{v}\ _2 < 2$ then it is possible that $\ \mathbf{u} - \mathbf{v}\ _2 > 2.5$ for certain values of $\mathbf{u}, \mathbf{v}$ .	
6	Classification models such as SVMs are more confident about their predictions on data points that lie very close to their decision boundary.	

**Q2. (Sliding parabolas)** Consider the two functions  $f(x) = (x - a)^2 + b$  and  $g(x) = x^4 + x^2$ .

Find values of $a, b \in \mathbb{R}$ such that the functions $f, g$ share a tangent at $x = 1$ .	$a =$	$b =$
For the values you find above, find out the value of the function $f$ at $x = 1$ .	$f(1) =$	
Write down the equation of the (shared) tangent of the functions at $x = 1$ .		
Find values of $a, b \in \mathbb{R}$ such that the functions $f, g$ share a stationary point $x_0$ such that $f(x_0) = g(x_0)$ .	$a =$	$b =$

Note that your equation for the tangent must not contain variables like  $a, b$  and should contain only absolute constants. Write your answers only in the space provided. **(2 + 1 + 1 + 2 = 6 marks)**

**Q3 (Robust regression)** Training with the absolute loss can make regression models less sensitive to outliers. Let us analyse a simplified version of the objective function used in this technique.

- a. Consider the function  $f(x) = |b \cdot x - a|$  where  $a, b, x \in \mathbb{R}$ . For any value of  $a, b, x_0 \in \mathbb{R}$ , write down an expression for the (entire) subdifferential of  $f$  at  $x_0$ . No derivation needed.

- b. Consider the function  $g(\mathbf{w}) = \frac{1}{2} \|\mathbf{w}\|_2^2 + |\mathbf{w}^\top \mathbf{x} - y|$  where  $\mathbf{w}, \mathbf{x} \in \mathbb{R}^d$  and  $y \in \mathbb{R}$ . Write down an expression for the (entire) subdifferential of  $g$  at  $\mathbf{w}^0$  for any value of  $\mathbf{w}^0, \mathbf{x} \in \mathbb{R}^d$  and  $y \in \mathbb{R}$ . No derivation needed. Write your answer in the space given. **(2 + 2 = 4 marks)**

Your answer to part a.

Your answer to part b.

**Q4. (Vector line-up)** Give examples of 4D vectors (fill-in the 4 boxes) with the following properties. Any example will get full marks so long as it satisfies the property mentioned in the question part. Your answers to the parts a, b, c, d may be same/different. **(4 x 1 = 4 marks)**

- a. A vector  $\mathbf{v} \in \mathbb{R}^4$  such that  $\mathbf{v} \neq \mathbf{0}$  and  $\mathbf{v}^\top \mathbf{u} = 0$  where  $\mathbf{u} = (1, -1, 1, -1) \in \mathbb{R}^4$ .
- b. A vector  $\mathbf{v} \in \mathbb{R}^4$  with unit  $L_3$  norm i.e.,  $\|\mathbf{v}\|_3 = 1$ .
- c. A vector  $\mathbf{v} \in \mathbb{R}^4$  with  $\|\mathbf{v}\|_2 \leq 1$  and  $\|\mathbf{v} - \mathbf{u}\|_2 \geq 2$  where  $\mathbf{u} = \left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right) \in \mathbb{R}^4$ .
- d. A vector  $\mathbf{v} \in \mathbb{R}^4$  such that  $\mathbf{v} \neq \mathbf{0}$  and whose  $L_2$  norm is half its  $L_1$  norm i.e.,  $\|\mathbf{v}\|_2 = \frac{1}{2} \|\mathbf{v}\|_1$ .
